

FIG. 1-1

1/31

Constitutively Active Receptors

File Name	Receptor	Mutation Site	Sequence	Assay / Cells	Reference
CLASS A GROUP I					
MSHR_mouse	melanocyte-stimulating hormone	TMII	92 <u>VSIIVLERTIL</u> SEQ ID NO: 2 K	adenylyl cyclase activity/ HEK293, stably transfected	(Robbins, Nadeau et al. 1993)
MSH					
CLASS A GROUP II					
SH1B_human	5-hydroxytryptamine _{1B}	C-terminus of IC3	313 RERKATKTLGI SEQ ID NO: 3 K, R, Q	binding of [³ S]GTP[S] / CHO-KJ	(Pauwels, Gouble et al. 1999)
SH2A_human	5-hydroxytryptamine _{2A}	C-terminus of IC3	322 NEQQKACKVVLGI SEQ ID NO: 4 K	IP production / COS-7	(Egan, Herrick-Davis et al. 1998)
2H2C_rat	5-hydroxytryptamine _{2C}	C-terminus of IC3	312 NEDDAS <u>SKVVLGI</u> SEQ ID NO: 5 L	PI hydrolysis / COS-7	(Herrick-Davis, Egan et al. 1997)

FIG. 1-2

2/31

CLASS A GROUP II					
A1AB_human	α_{1B} -adrenergic alpha 1B-AR	TMDI junction between TMDIII and IC2	FAIVGNILVIL SEQ ID NO: 6 A 142 CAISIDRYIGV SEQ ID NO: 7 A	IP / COS-7	(Scheer, Famili et al. 1997)
A1ABB_human	α_{1B} -adrenergic alpha 1B-AR	TMDII junction between TMDIII and IC2	CAISIDRYIGV SEQ ID NO: 8 K	IP / COS-7	(Scheer, Costa et al. 2000)
A1ABBHuman	α_{1B} -adrenergic α_{1B} -adrenergic	TMIII carboxyl end of IC3 TMV	AVDVLQCTASI SEQ ID NO: 9 F 293 REKKAAKTLGI SEQ ID NO: 10 E 204 EEPFPYALFSSILG SEQ ID NO: 11 V	IP / COS-1 arachidonic acid release IP / COS-1	(Perez, Hwa et al. 1996) (Hwa, Gaivin et al. 1997)
A1ABHuman	α_{1B} -adrenergic	C-terminal IC3	SRERKAAKT SEQ ID NO: 12 X=19 different substitutions	PI / COS-7	(Kjelsberg, Cotecchia et al. 1992)
A1ABHuman	α_{1B} -adrenergic	C-terminal IC3	KFSSEKKAAKTGLGI SEQ ID NO: 13 K H L	PI hydrolysis / rat fibroblast	(Allen, Leikowitz et al. 1991)
A2AAHuman	α_2 C10-adrenergic alpha-2AAR	C-terminal IC3 loop	273 (348?) EKRTFTVLA X=F, A, C, E, K	Adenyl cyclase inhibition / HEK293	(Ren, Kurose et al. 1993)
ACM1Human	muscarinic Hm1	C-terminal IC3 loop junction	360 SLVREKKAARTLS A	PI / HEK(U23)	(Högger, Shocket et al. 1995)
ACM2-human	muscarinic acetylcholine M1 muscarinic acetylcholine M2	junction of IC3 and TMVI	390 KRYVPTIL'A 1-4 A inserted	IP production, inhibition of cAMP production / COS-7	(Liu, Blin et al. 1996)

FIG. 1-3

3/31

CLASS A GROUP II		TMVI			
ACM3_rat	m3 muscarinic (rat) muscarnic acetylcholine M3	TMVI	507 TWT ^S PYNNIMVLVNT SEQ ID NO: 17	IP / COS-7	(Blüml, Mutschler et al. 1994)
ACM5_human	m5 muscarinic muscarnic acetylcholine M5	N-terminus to TMII TMVI	chimera composed of m2 I-69 m5 77-445 m2 391-466	β-gal / NIH 3T3	(Burstein, Spalding et al. 1996)
ACM5_human	m5 muscarinic muscarnic acetylcholine M5	TMVI	451 A ^M LLA E ^L ITW T ^V PYNI M ^H VLY ^S T V V S F T	β-gal; radioligand binding / NIH-3T3	(Spalding, Burstein et al. 1998)
ACM5_humman	m5 muscarinic muscarnic acetylcholine M5	junction of TMVI and EC3	465 Y ^X NIMLV ^S TFCDDKCV SEQ ID NO: 19 X=V,F,R,K,+more	β-gal; radioligand binding / NIH-3T3	(Spalding, Burstein et al. 1997)
B1AR_human	β ₁ -adrenergic	C-terminus	389 RKA ^R FQGLLCCA SEQ ID NO: 20	adenylyl cyclase; agonist binding / CHW	(Mason, Moore et al. 1999)
B2AR_human	β ₂ -adrenergic beta-2AR	C-terminal IC3 loop	266 272 FCL ^K EHHK ^A LKT ^I LGI SEQ ID NO: 21 SR K A	adenylyl cyclase activation; agonist binding affinity / COS-7 or CHO	(Samama, Cotecchia et al. 1993); (Lefkowitz, Cotecchia et al. 1993)
DADR_human	dopamine D1A	carboxyl terminal IC3	264 SFKMSEKRET ^I KVLKT SEQ ID NO: 22 K 288 from D1B receptor APDTSIKKET ^K VLKT SEQ ID NO: 23	adenylyl cyclase; cAMP accumulation / HEK293	(Charpentier, Jarvie et al. 1996)
DADR_human	dopamine D1	TMVI	286 FVCCW ^N LPPFFIL SEQ ID NO: 24 A	cAMP accumulation / COS-7	(Cho, Taylor et al. 1996)
HH2R_rat	histamine H ₂	IC2	115 FMISLD ^N RYCAV SEQ ID NO: 25 A	cAMP production / HEK-293	(Alewijnse, Timmerman et al. 2000)

FIG. 1-4

4/31

File Name	Receptor	Mutation Site	Sequence	Assay / Cells	Reference
CLASS A GROUP III					
OPSD_human	opsin rhodopsin	TMII	90 D 113 GCNLEGGFFAT Q 292 296	transducin; phosphorylation by rhodopsin kinase / COS	(Rim and Oprian 1995)
		TMIII	MTIPAFPA <u>K</u> SAAIY SEQ ID NO: 28 E, G, E, M		
		TMVII	²⁹² Ala neutral a.a converted to carboxylate and competes with ¹¹³ Glu for salt bridge with ²⁹⁶ Lys		
OPSD_human	opsin rhodopsin	TMII	134 VVLA <u>I</u> E <u>R</u> YVVV SEQ ID NO: 29 I, Q, S	transducin; radioligand binding / COS	(Acharya and Karnik 1996)
OPSD_human	opsin rhodopsin	TM6	257 RMVIIIMVIAFL SEQ ID NO: 30 Y, N	transducin, GTP _γ S uptake / COS	(Han, Smith et al. 1998)
OPSD_human	opsin rhodopsin	plus TM3 TMVII	plus G113Q 296 PAFF <u>A</u> K <u>S</u> AAIY SEQ ID NO: 31 G X=E,M natural mutants + 10 different a.a. substitutions	transducin; radioligand binding / COS	(Govardhan and Oprian 1994); (Cohen, Yang et al. 1993)
OPSD_human	opsin rhodopsin	IC2	134 VVLA <u>I</u> E <u>R</u> YVVV SEQ ID NO: 32 Q	disrupts critical salt bridge between ²⁹⁶ Lys(TMVII) and ¹¹³ Glu(TMIII)	(Cohen, Yang et al. 1993)

FIG. 1-5

5/31

TRFR_mouse	thyrotropin-releasing hormone TRH-R	carboxyl tail	335 FRKLCNCQK STOP	SEQ ID NO: 33	⁴ Ca ²⁺ efflux, [Ca ²⁺] Xenopus oocytes; IP formation / ArT20, <i>stably transfected</i>	(Matus-Leibovitch, Nussenzveig et al. 1995)
------------	--	---------------	--------------------------	---------------	---	--

FIG. 1-6

6/31

File Name	Receptor	Mutation Site	Sequence	Assay / Cells	Reference
CLASS A GROUP IV BRB2_human	bradykinin B ₂	TMIII	AIISMNLYSSI A	IP production / COS-7	(Marie, Koch et al. 1999)
	B2 bradykinin BK-2	TMVI	256 LLFIICWLPFQI F	SEQ ID NO: 34 SEQ ID NO: 35	

FIG. 1-7

7/31

File Name	Receptor	Mutation Site	Sequence	Assay / Cells	Reference
CLASS A GROUP V					
AG2R_rat	AT _{1A}	TMIII	111 ASVSFILYASV SEQ ID NO: 36 A	phospholipase C; IP production / COS-7	(Grobleski, Maignet et al. 1997)
AG2R_rat	Type-1A angiotensin II	C-terminus of TM7 other multiple mutations	305 LFYGF _Q GKKFK SEQ ID NO: 37	IP production / HEK- 293; intracellular Ca ²⁺ mobilization / CHO	(Parnot, Bardin et al. 2000)
FMLR_human	formylmethionylleucylphenylal anine (fMLPR)	IC1	51 LVIWV _E AGPFMTHT _T ISYLNKA SEQ ID NO: 38 SEQ ID NO: 39	PI production; phospholipase C stimulation / COS-7	(Amatuda, Draga- Graonic et al. 1995)
IL8B_human	interleukin-8 receptor B CXCR-2 chemokine	IC2	138 ACISV _V RYLAIVH SEQ ID NO: 40	IP production; Ca ²⁺ mobilization and actin polymerization / NIH 3T3	(Burger, Burger et al. 1999)
LSHR_human	luteinizing hormone (LH)	IC3	564 MATN _G TKIAKK SEQ ID NO: 41	cAMP production / HEK293	(Kudo, Osuga et al. 1996)
LSHR_human	luteinizing hormone (LH)	TMVI	578 ILLIF _G DFTCM SEQ ID NO: 42	cAMP production / COS-7	(Shenker, Lau et al. 1993)
LSHR_human	luteinizing hormone (LH)	TM6	571 KIAK _I KI _I LLIFTDFTCM SEQ ID NO: 43	cAMP production / COS-7	(Kosugi, Van Dop et al. 1995)
LSHR_rat	luteinizing hormone / human chorionic gonadotropin (LH/hCG)	TMVI	556 ILLIF _G DFTCM SEQ ID NO: 44	cAMP production / HEK 293T	(Bradbury, Kawate et al. 1997; Bradbury and Meron 1999)
OPRD_mouse	delta opioid receptor	TM3	128 KVLS _{A,X,H} YYDMF SEQ ID NO: 45	adenylyl cyclase inhibition / COS-7	(Cavalli, Babey et al. 1999)
OXYR_human	oxytocin	IC2	137 LMSLD _A GLAIC SEQ ID NO: 46	IP production / COS-7	(Fanelli, Barbier et al. 1999)

FIG. 1-8

8/31

PAFR_human	platelet-activating factor (PAF)	C-terminus of IC3	231 EVKRRALWMVCTLVAV SEQ ID NO: 47 R	IP production / COS-7	(Parent, Le Gouill et al. 1996)
PAFR_human	platelet-activating factor (PAF)	TMIII	100 CLFFINTYCSV SEQ ID NO: 48 A	arachidonate release, IP production, adenylyl cyclase inhibition / CHO	(Ishii, Izumi et al. 1997)
PE23_human	prostaglandin E ₁ , E13II E13IV	C-terminal tail	360 FCQEEFWGN SEQ ID NO: 49 FCQMRKRLREQQEFWGN SEQ ID NO: 50 ↑ truncated	inhibition of adenylyl cyclase / CHO, stably expressed	(Jin, Mao et al. 1997)
PE23_mouse	prostaglandin E ₁ , EP3	carboxyl-terminal tail SEQ ID NO: 51	336 KILLRKFCQIRDHT (3α) MMNNHL (3β) ↑ truncated	inhibition of adenylyl cyclase / CHO, stably expressed	(Hasegawa, Negishi et al. 1996)
THTRHuman	thrombin	EC2 loop SEQ ID NO: 52	259 CHDVLNNETLLEGYYAYY DLKD KDF I F, M	“Ca ²⁺ efflux, PI hydrolysis, reporter gene induction / COS-7	(Nanevitz, Wang et al. 1996)
TSHRHuman	thyrotropin (TSHR) thyroid stimulating hormone	EC1 EC2	486 YYNHADWQTG SEQ ID NO: 53 YAKVSICLPMD SEQ ID NO: 54 F, M T	inositol phosphate-- diacylglycerol cascade / COS-7	(Parma, Van Sande et al. 1995)
TSHRHuman	thyrotropin (TSHR) thyroid stimulating hormone	TMIII TMVII	509 ASELSVYTLTV SEQ ID NO: 55 A 672 YPLNSCAMPPFL SEQ ID NO: 56 Y	adenylyl cyclase activation / COS-7	(Duprez, Parma et al. 1994)
TSHRHuman	thyrotropin (TSHR) thyroid stimulating hormone	TMV	597 VAFVIVCCCHV SEQ ID NO: 57 L	cAMP formation / COS-7 cells	(Esapa, Duprez et al. 1999)
TSHRHuman	thyrotropin (TSHR)	TMVII	677 CANPFLYAIFT SEQ ID NO: 58 V	cAMP formation / CHO cells	(Russo, Wong et al. 1999)
TSHRHuman	thyrotropin (TSHR) thyroid stimulating hormone	IC3	613 VRNPQYNPGDKTKIAK deletion SEQ ID NO: 59	cAMP formation / COS-7	(Wonerow, Schoneberg et al. 1998)

FIG. 1-9

9/31

TSHR_human	thyrotropin (TSHR) thyroid stimulating hormone	IC3 / TMVI SEQ ID NO: 60	623 KDTKIAKRMAVLIFTDFICM V I	cAMP activation / COS-7	(Paschke, Tonacchera et al. 1994)
V2R_human	vasopressin V ₂	IC2 SEQ ID NO: 61	136 LAMTLIDHRRAI A	cAMP formation / COS-7	(Morin, Cotte et al. 1998)

FIG. 1-10

10/31

File Name	Receptor	Mutation Site	Sequence	Assay / Cells	Reference
CLASS B GROUP I					
CALR_human	human calcitonin hCTR-1 hCTR-2	wild type (native) protein			(Cohen, Thaw et al. 1997)
CLASS B GROUP II					
PTRR_human	parathyroid hormone PTH / PTH-related peptide	junction of IC1 and TMII	223 TRNYIHMHLFL SEQ ID NO: 62 R, K	cAMP accumulation / COS-7	(Schipani, Jensen et al. 1997)
		junction of IC3 and TMVI	410 KLLKSTLVLM P SEQ ID NO: 63 C, others		
CLASS B GROUP III					
GIPR_human	Glucose-dependent insulinotropic peptide (GIP-R)	TMVI	340 VFAPVTEEQAR SEQ ID NO: 64 P	cAMP production / L293	(Tseng and Lin 1997)
GLR_rat	glucagon	junction of IC1 loop 1 and TMII	178 TRNYIHGNLFA SEQ ID NO: 65 R	cAMP accumulation / COS-7	(Hjorth, Orskov et al. 1998)
		IC end of TMVI	352 RLARSTLTLIP SEQ ID NO: 66 A		
VIPR_human	vasoactive intestinal peptide 1 (VIP)	junction of IC loop 1 and TMII	178 RNYIHMHLFI SEQ ID NO: 67 R	cAMP production / COS-7 or CHO	(Gaudin, Maoret et al. 1998)
		junction of IC loop 3 and TMVI	343 LARSTLLIP SEQ ID NO: 68 X= K, P		(Gaudin, Rouyer-Fessard et al. 1998)

FIG. 1-11

11/31

File Name	Receptor	Mutation Site	Sequence	Assay / Cells	Reference
CLASS_C					
CASR_human	calcium-sensing	N-terminal EC	TLSFVAQN KIDS LDEFNCSEH I	IP / IsA	(Jensen, Spalding et al. 2000)
			various substitutions, in multiple combinations	SEQ ID NO: 69	

FIG. 1-12

12/31

File Name	Receptor	Mutation Site	Sequence	Assay / Cells	Reference
CLASS_D					
O74283 RCB2 <i>C. cinereus</i>	pheromone	TM6	229 PLSAYQIYILGR SEQ ID NO: 70 P	heterologous yeast assay	(Olesnicki, Brown et al. 1999)
STE2_yeast	pheromone α -factor	TM6	258 QSLL VPSIIIFI SEQ ID NO: 71 L,L	<i>lacZ</i> reporter gene	(Konopka, Margarit et al. 1996)
STE2_yeast	pheromone α -factor	double mutations TM5 and TM6	223 MSPV L VVK I LAIR SEQ ID NO: 72 C C 247 251 DSF HILL I LCQSLL SEQ ID NO: 73 CC CC	<i>lacZ</i> reporter gene / yeast	(Dube, DeCostanzo et al. 2010)
STE3_yeast	pheromone α -factor	IC3	194 DVRD I L HCTNS SEQ ID NO: 74 Q	β -galactosidase	(Boone, Davis et al. 1993)
STE2_yeast	pheromone α -factor	TM6	253 258 LIMSCQ SLLVPSIIIFI SEQ ID NO: 75 L L P	β -galactosidase	(Sommers, Martin et al. 2000)

FIG. 1-13

Bibliography

13/31

Acharya, S. and S. S. Karnik (1996). "Modulation of GDP release from transducin by the conserved Glu134-Arg135 sequence in rhodopsin." *J Biol Chem* 271(4): 25406-11.

Alewijne, A. E., H. Timmerman, et al. (2000). "The Effect of Mutations in the DRY Motif on the Constitutive Activity and Structural Instability of the Histamine H(2) Receptor." *Mol Pharmacol* 57(5): 890-898.

Allen, L. F., R. J. LeRowitz, et al. (1991). "G-protein-coupled receptor genes as protooncogenes: constitutively activating mutation of the alpha 1B-adrenergic receptor enhances mitogenesis and tumorigenicity." *Proc Natl Acad Sci U S A* 88(24): 11354-8.

Amatruda, T. T., 3rd, S. Dragas-Graonic, et al. (1995). "Signal transduction by the formyl peptide receptor. Studies using chimeric receptors and site-directed mutagenesis define a novel domain for interaction with G-proteins." *J Biol Chem* 270(47): 28010-3.

Büuml, K., E. Mutschler, et al. (1994). "Functional role in ligand binding and receptor activation of an asparagine residue present in the sixth transmembrane domain of all muscarinic acetylcholine receptors." *J Biol Chem* 269(29): 18870-6.

Boone, C., N. G. Davis, et al. (1993). "Mutations that alter the third cytoplasmic loop of the a-factor receptor lead to a constitutive and hypersensitive phenotype." *Proc Natl Acad Sci U S A* 90(21): 9921-5.

Bradbury, F. A., N. Kawate, et al. (1997). "Post-translational processing in the Golgi plays a critical role in the trafficking of the luteinizing hormone/human chorionic gonadotropin receptor to the cell surface." *J Biol Chem* 272(9): 5921-6.

Bradbury, F. A. and K. M. Menon (1999). "Evidence that constitutively active luteinizing hormone/human chorionic gonadotropin receptors are rapidly internalized." *Biochemistry* 38(27): 8703-12.

Burger, M., J. A. Burger, et al. (1999). "Point mutation causing constitutive signaling of CXCR2 leads to transforming activity similar to Kaposi's sarcoma herpesvirus-G protein-coupled receptor." *J Immunol* 163(4): 2017-22.

Burstein, E. S., T. A. Spalding, et al. (1996). "Constitutive activation of chimeric m2/m5 muscarinic receptors and delineation of G-protein coupling selectivity domains." *Biochem Pharmacol* 51(4): 539-44.

Cavalli, A., A. M. Babey, et al. (1999). "Altered adenylyl cyclase responsiveness subsequent to point mutations of Asp 128 in the third transmembrane domain of the delta-opioid receptor." *Neuroscience* 93(3): 1025-31.

Charpentier, S., K. R. Larvie, et al. (1996). "Silencing of the constitutive activity of the dopamine D1B receptor. Reciprocal mutations between D1 receptor subtypes delineate residues underlying activation properties." *J Biol Chem* 271(45): 28071-6.

Cho, W., L. P. Taylor, et al. (1996). "Mutagenesis of residues adjacent to transmembrane prolines alters D1 dopamine receptor binding and signal transduction." *Mol Pharmacol* 50(5): 1338-45.

Cohen, D. P., C. N. Thaw, et al. (1997). "Human calcitonin receptors exhibit agonist-independent (constitutive) signaling activity." *Endocrinology* 138(4): 1400-5.

Cohen, G. B., T. Yang, et al. (1993). "Constitutive activation of opsin: influence of charge at position 134 and size at position 296." *Biochemistry* 32(23): 6111-5.

Dube, P., A. DeCostanzo, et al. (2000). "Interaction between transmembrane domains five and six of the alpha -factor receptor." *J Biol Chem* 275(34): 26492-9.

Duprez, L., J. Parma, et al. (1994). "Germline mutations in the thyrotropin receptor gene cause non- autoimmune dominant hyperthyroidism." *Nat Genet* 7(3): 396-401.

Egan, C. T., K. Herrick-Davis, et al. (1998). "Creation of a constitutively activated state of the 5- hydroxytryptamine2A receptor by site-directed mutagenesis: inverse agonist activity of antipsychotic drugs." *J Pharmacol Exp Ther* 286(1): 85-90.

Esapa, C. T., L. Duprez, et al. (1999). "A novel thyrotropin receptor mutation in an infant with severe thyrotoxicosis." *Thyroid* 9(10): 1005-10.

Fanelli, F., P. Barbier, et al. (1999). "Activation mechanism of human oxytocin receptor: a combined study of experimental and computer-simulated mutagenesis." *Mol Pharmacol* 56(1): 214-25.

Gaudin, P., J. J. Madret, et al. (1998). "Constitutive activation of the human vasoactive intestinal peptide 1 receptor, a member of the new class II family of G protein-coupled receptors." *J Biol Chem* 273(9): 4990-6.

Gaudin, P., C. Rouyer-Fessard, et al. (1998). "Constitutive activation of the human VIP1 receptor." *Annu NY Acad Sci* 865: 382-5.

FIG. 1-14

Govardhan, C. P. and D. D. Oprian (1994). "Active site-directed inactivation of constitutively active mutants of rhodopsin." *J Biol Chem* 269(9): 6524-7.

Groblewski, T., B. Maigret, et al. (1997). "Mutation of Asn111 in the third transmembrane domain of the AT1A angiotensin II receptor induces its constitutive activation." *J Biol Chem* 272(3): 1822-6.

Han, M., S. O. Smith, et al. (1998). "Constitutive activation of opsin by mutation of methionine 257 on transmembrane helix 6." *Biochemistry* 37(22): 8253-61.

Hassegawa, H., M. Negishi, et al. (1996). "Two isoforms of the prostaglandin E receptor EP3 subtype different in agonist-independent constitutive activity." *J Biol Chem* 271(4): 1857-60.

Herrick-Davis, K., C. Egan, et al. (1997). "Activating mutations of the serotonin 5-HT2C receptor." *J Neurochem* 69(3): 1138-44.

Hjorth, S. A., C. Orskov, et al. (1998). "Constitutive activity of glucagon receptor mutants." *Mol Endocrinol* 12(1): 78-86.

Högger, P., M. S. Shockley, et al. (1995). "Activating and inactivating mutations in N- and C-terminal i3 loop junctions of muscarinic acetylcholine M1 receptors." *J Biol Chem* 270(13): 7405-10.

Hwa, J., R. Graivin, et al. (1997). "Synergism of constitutive activity in alpha 1-adrenergic receptor activation." *Biochemistry* 36(3): 633-9.

Ishii, I., T. Izumi, et al. (1997). "Alanine exchanges of polar amino acids in the transmembrane domains of a platelet-activating factor receptor generate both constitutively active and inactive mutants." *J Biol Chem* 272(12): 7846-54.

Jensen, A. A., T. A. Spalding, et al. (2000). "Functional importance of the Ala116-Pro136 region in the calcium-sensing receptor. CONSTITUTIVE ACTIVITY AND INVERSE AGONISM IN A FAMILY C G-PROTEIN-COUPLED RECEPTOR [In Process Citation]." *J Biol Chem* 275(38): 29547-55.

Jin, J., G. F. Mao, et al. (1997). "Constitutive activity of human prostaglandin E receptor EP3 isoforms." *British J Pharmacol* 121: 317-23.

Kjelsberg, M. A., S. Cotechia, et al. (1992). "Constitutive activation of the alpha 1B-adrenergic receptor by all amino acid substitutions at a single site. Evidence for a region which constrains receptor activation." *J Biol Chem* 267(3): 1430-3.

Konopka, J. B., S. M. Margarini, et al. (1996). "Mutation of Pro-258 in transmembrane domain 6 constitutively activates the G protein-coupled alpha-factor receptor." *Proc Natl Acad Sci U S A* 93(13): 6764-9.

Kosugi, S., C. Van Dop, et al. (1995). "Characterization of heterogeneous mutations causing constitutive activation of the luteinizing hormone receptor in familial male precocious puberty." *Hum Mol Genet* 4(2): 183-8.

Kudo, M., Y. Osuga, et al. (1996). "Transmembrane regions V and VI of the human luteinizing hormone receptor are required for constitutive activation by a mutation in the third intracellular loop." *J Biol Chem* 271(37): 22470-8.

Leikowitz, R. J., S. Cotechia, et al. (1993). "Constitutive activity of receptors coupled to guanine nucleotide regulatory proteins." *Trends Pharmacol Sci* 14(8): 303-7.

Liu, J., N. Blin, et al. (1996). "Molecular mechanisms involved in muscarinic acetylcholine receptor-mediated G protein activation studied by insertion mutagenesis." *J Biol Chem* 271(11): 6172-8.

Marie, J., C. Koch, et al. (1999). "Constitutive activation of the human bradykinin B2 receptor induced by mutations in transmembrane helices III and VI." *Mol Pharmacol* 55(1): 92-101.

Mason, D. A., J. D. Moore, et al. (1999). "A gain-of-function polymorphism in a G-protein coupling domain of the human beta1-adrenergic receptor." *J Biol Chem* 274(18): 12670-4.

Manus-Leibovitch, N., D. R. Nussenzveig, et al. (1995). "Truncation of the thyrotropin-releasing hormone receptor carboxyl tail causes constitutive activity and leads to impaired responsiveness in Xenopus oocytes and AT20 cells." *J Biol Chem* 270(3): 1041-7.

Morin, D., N. Corre, et al. (1998). "The D136A mutation of the V2 vasopressin receptor induces a constitutive activity which permits discrimination between antagonists with partial agonist and inverse agonist activities." *Eur J Biochem* 241(3): 470-5.

Natevicz, T., L. Wang, et al. (1996). "Thrombin receptor activating mutations. Alteration of an extracellular agonist recognition domain causes constitutive signaling." *J Biol Chem* 271(2): 702-6.

Olesnicky, N. S., A. J. Brown, et al. (1999). "A constitutively active G-protein-coupled receptor causes mating self-compatibility in the mushroom Coprinus." *Embo J* 18(10): 2756-63.

PARENT, J. L., C. LE GOUIL, et al. (1996). "Mutations of two adjacent amino acids generate inactive and constitutively active forms of the human platelet-activating factor receptor." *J Biol Chem* 271(14): 7949-55.

14/31

FIG. 1-15

Parma, J., J. Van Sande, et al. (1995). "Somatic mutations causing constitutive activity of the thyrotropin receptor are the major cause of hyperfunctioning thyroid adenomas: identification of additional mutations activating both the cyclic adenosine 3',5'-monophosphate and inositol phosphate-Ca²⁺ cascades." *Mol Endocrinol* 9(6): 725-33.

Parnot, C., S. Bardin, et al. (2000). "Systematic identification of mutations that constitutively activate the angiotensin II type 1A receptor by screening a randomly mutated cDNA library with an original pharmacological bioassay." *Proc Natl Acad Sci U S A* 97(13): 7615-20.

Paschke, R., M. Tonacchera, et al. (1994). "Identification and functional characterization of two new somatic mutations causing constitutive activation of the thyrotropin receptor in hyperfunctioning autonomous adenomas of the thyroid." *J Clin Endocrinol Metab* 139(6): 1785-9.

Pauwels, P. J., A. Gouble, et al. (1999). "Activation of constitutive 5-hydroxytryptamine1B receptor by a series of mutations in the BBXXB motif: positioning of the third intracellular loop distal junction and its goalpha protein interactions [In Process Citation]." *Biochem J* 343 Pt 2: 435-42.

Perez, D. M., J. Hwa, et al. (1996). "Constitutive activation of a single effector pathway: evidence for multiple activation states of a G protein-coupled receptor." *Mol Pharmacol* 49(1): 112-22.

Ren, Q., H. Kurose, et al. (1993). "Constitutively active mutants of the alpha 2-adrenergic receptor [published erratum appears in *J Biol Chem* 1994 Jan 14;269(2):1566]." *J Biol Chem* 268(22): 16483-7.

Kim, J. and D. D. Oprian (1995). "Constitutive activation of opsin: interaction of mutants with rhodopsin kinase and arrestin." *Biochemistry* 34(37): 11938-45.

Robbins, L. S., J. H. Nadeau, et al. (1993). "Pigmentation phenotypes of variant extension locus alleles result from point mutations that alter MSH receptor function." *Cell* 72(6): 827-34.

Russo, D., M. G. Wong, et al. (1999). "A Val 677 activating mutation of the thyrotropin receptor in a Hurthle cell thyroid carcinoma associated with thyrotoxicosis." *Thyroid* 9(1): 13-7.

Sartama, P., S. Cotecchia, et al. (1993). "A mutation-induced activated state of the beta 2-adrenergic receptor. Extending the ternary complex model." *Journal of Biological Chemistry* 268(7): 4625-36.

Scheer, A., T. Costa, et al. (2000). "Mutational analysis of the highly conserved arginine within the Glu/Asp-Arg-Tyr motif of the alpha(1b)-adrenergic receptor: effects on receptor isomerization and activation." *Mol Pharmacol* 57(2): 219-31.

Scheer, A., F. Fanelli, et al. (1997). "The activation process of the alpha1B-adrenergic receptor: potential role of protonation and hydrophobicity of a highly conserved aspartate." *Proc Natl Acad Sci U S A* 94(3): 808-13.

Schipani, E., G. S. Jensen, et al. (1997). "Constitutive activation of the cyclic adenosine 3',5'-monophosphate signaling pathway by parathyroid hormone (PTH)/PTH-related peptide receptors mutated at the two loci for Jansen's metaphyseal chondrodyplasia." *Mol Endocrinol* 11(7): 851-8.

Shenker, A., L. Laue, et al. (1993). "A constitutively activating mutation of the luteinizing hormone receptor in familial male precocious puberty [see comments]." *Nature* 365(6447): 652-4.

Sommers, C. M., N. P. Martin, et al. (2000). "A limited spectrum of mutations causes constitutive activation of the yeast alpha-factor receptor." *Biochemistry* 39(23): 6898-909.

Spalding, T. A., E. S. Burstein, et al. (1998). "Identification of a ligand-dependent switch within a muscarinic receptor." *J Biol Chem* 273(34): 21563-8.

Spalding, T. A., E. S. Burstein, et al. (1997). "Constitutive activation of the m5 muscarinic receptor by a series of mutations at the extracellular end of transmembrane 6." *Biochemistry* 36(33): 10109-16.

Tseng, C. C. and L. Lin (1997). "A point mutation in the glucose-dependent insulinotropic peptide receptor confers constitutive activity." *Biochem Biophys Res Commun* 233(1): 96-100.

Wonenow, P., T. Schoneberg, et al. (1998). "Deletions in the third intracellular loop of the thyrotropin receptor. A new mechanism for constitutive activation." *J Biol Chem* 273(14): 7900-5.

15/31

FIG. 2

A Point Mutation Enhances MC-4 Receptor
Constitutive Activity




FIG. 3

Light Emission Induced by the WT CCK-BR
vs. a Constitutively Active Mutant

17/31




FIG. 4

A Point Mutation Confers Constitutive Activity
to the Rat μ Opioid Receptor




FIG. 5

Forskolin Stimulated HEK293 Cells Transfected
With pcDNA1 and a CRE-luc Construct




FIG. 6

The Rat μ Opioid Receptor Signals Through G α i




FIG. 7

A Point Mutation Confers Constitutive Activity
to the Rat μ Opioid Receptor




FIG. 8

Target Residues Within Class I GPCRs




FIG. 9
TMD III Asn (-14 from DRY) is a Target
for Mutation Induced Constitutive Activity





FIG. 10

The 'DRY' Motif is a Target for Mutation
Induced Constitutive Activity



24/31

FIG. 11

A Point Mutation Enhances MC-4 Receptor Constitutive Activity





FIG. 12

The -13 Position is a Target for Mutation
Induced Constitutive Activity



27/31

FIG. 13

SEQ ID NO: 76	ork	1	-----MESPIQIFRGEPGPTCAPSACLPPNSAWFPGWAEP..DSNGSAGCSEDAC
SEQ ID NO: 77	orkr	1	-----MESPIQIFRGEPGPTCAPSACLPPNSSWFPNWAES..DSNGSVCSEDQQ
SEQ ID NO: 78	orm	1	MDSSAAPTNASNCTDALAYSSCSPAPSEGSWV..NLSHLDGNLSDPCGPNRDLDLGGRDSDL
SEQ ID NO: 79	ormr	1	MDSSTGPGNTSDCSDPQAQASCSPA..PGSWL..NLSHVDGNQSDPCGLNRGLGGNDSDL
SEQ ID NO: 80	ord	1	-----MEPAPSAGAE..Q.PPLFTANASDAYPSACPSAGANASG
SEQ ID NO: 81	AT1a	1	-----MALNSSAEDGIKRIQ
SEQ ID NO: 82	BK-2	1	-----MFSPWKISMFLSVREDSVPTTASFSADMNVTLQGPTLNG.TFAQ
ork	49	LEPAEISPAF..PVVITAVYYSVVFVVGVLGNNSLVMEVITYRTKMKTTATNIYIFNLALADA	
orkr	49	LEPAEISPAF..PVVITAVYYSVVFVVGVLGNNSLVMEVITYRTKMKTTATNIYIFNLALADA	
orm	59	CPTGTS.PSMVTAITIMALYSIVCVVGLFGNFLVMVIVIYRTKMKTTATNIYIFNLALADA	
ormr	57	CPTGTS.PSMVTAITIMALYSIVCVVGLFGNFLVMVIVIYRTKMKTTATNIYIFNLALADA	
ord	37	PPGANSASSALALAITALYSAVCAGLIGGNVLVMEGIVIYRTKMKTTATNIYIFNLALADA	
AT1a	16	DDCPKAAGRHSYIYVMIPTLYSIVFVVGIFGNSLVIVIVYFMKIKIVASVFLINLALADL	
BK-2	45	SKCPQVEWLWLNTIOPPFILWVLFVILATLENIFVIVSIFCLHKSSCTVAELYLGNLAADL	
ork	107	IYVHIMPFOSTVYLMN.SWPFGDVLCKIVISIDYYNMFTSIFTLTMSVDRYIAVCHPVK	
orkr	107	IYVHIMPFOSTVYLMN.SWPFGDVLCKIVISIDYYNMFTSIFTLTMSVDRYIAVCHPVK	
orm	118	LATSTLPFOSVNYLMG..TWPFGTILCKIVISIDYYNMFTSIFTLTMSVDRYIAVCHPVK	
ormr	116	LATSTLPFOSVNYLMG..TWPFGTILCKIVISIDYYNMFTSIFTLTMSVDRYIAVCHPVK	
ord	97	LATSTLPFOSVNYLMG..TWPFGTILCKIVISIDYYNMFTSIFTLTMSVDRYIAVCHPVK	
AT1a	76	CFLITLPLWVNTAMERYWPFGNHLCKIASASVTENLYASVFLICISIDRYIAVHPMK	
BK-2	105	ILACGLPEWLTISNNFDWLEGETILCRUUVNHIISMLY9SICFLMVSIDRYIALVKTM	
-14 from DRY * 			
ork	166	ALDFRTPLAKIIINICIWIASSVGISAIVIIGGTKVR..EDVDVIECSLOFEDDEYSWPD	
orkr	166	ALDFRTPLAKIIINICIWIASSVGISAIVIIGGTKVR..EDVDVIECSLOFEDDEYSWPD	
orm	177	ALDFRTPRNAKIIINYCNWLISSANGLPVMEATTKVR..C.GSIECTLIESHPTW.YWE	
ormr	175	ALDFRTPRNAKIIINYCNWLISSANGLPVMEATTKVR..C.GSIECTLIESHPTW.YWE	
ord	156	ALDFRTPAKAKIINICIWIASCVGVPIMVMAVTRPR..D.GAVVCMLOEESPSSW.YWD	
AT1a	136	SRLRRIMLVAKYTCIIWVAGLASIPEAVIHRNV..YFIENTNITVCAFHYESRN.STLP	
BK-2	165	MGRMRGVRWAKEYSIVIWCGLLISSEMIVFRTMKEYSDEGHNTACVISYES..LIWE	
ork	224	IFMKICVFIFAFVTPVLIITVCYTLMLIRLKSVRILSGSREKDRNLRRITRVLVVAVE	
orkr	224	IFMKICVFIFAFVTPVLIITVCYTLMLIRLKSVRILSGSREKDRNLRRITRVLVVAVE	
orm	232	NLKICVFIFAFIMPVLIITVCYGLMLIRLKSVRILSGSKEKDRNLRRITRVLVVAVE	
ormr	230	NLKICVFIFAFIMPVLIITVCYGLMLIRLKSVRILSGSKEKDRNLRRITRVLVVAVE	
ord	211	TVTKICVFIFAFVTPVLIITVCYGLMLIRLKSVRILSGSKEKDRSLRRITRVLVVAVE	
AT1a	193	IIGLGETKNILGFIFPFLITTSYTLIWKALKKAYEIQKPKRND..IFRLLIMATVLF	
BK-2	222	VFTNMLLNIVGFLIP.LSIVTFCTMQVQLRNNEQKFKEIQTE.RRAIVLVLVVAVE	
ork	284	IVCWTPIHIFIILVHALGS.T....SHSTAALSSYYFCIALGYTNSSLNPVLYAFLDENF	
orkr	284	IVCWTPIHIFIILVHALGS.T....SHSTAALSSYYFCIALGYTNSSLNPVLYAFLDENF	
orm	292	IVCWTPIHIFIILVHALGS.T....SHSTAALSSYYFCIALGYTNSSLNPVLYAFLDENF	
ormr	290	IVCWTPIHIFIILVHALGS.T....SHSTAALSSYYFCIALGYTNSSLNPVLYAFLDENF	
ord	271	IVCWAPIHIFIIVVWTLVDID....RRDPLVVAALHLCLIALGYANSSLNPVLYAFLDENF	
AT1a	250	FFSWVPHQIIFTFLDVLIQLGVIVHDCKISDIVDTAMPITICIAFYNNCLNPIFYGFLGKF	
BK-2	280	IVCWLPFOISTFLDTLHRIGILGILSSCQDERIUDVITQIASMAYNSCLNPVLYVIVGKF	
ork	338	KRCFRDFCFPLKMRMERQSTSRRP..NTVOD..BAYLRDIDGMNKPV-----	
orkr	338	KRCFRDFCFPLKMRMERQSTSRRP..NTVOD..BASMRDVGGMNKPV-----	
orm	346	KRCFRDFCFPLKMRMERQSTSRRP..NTVOD..BASMRDVGGMNKPV-----	
ormr	344	KRCFRDFCFPLKMRMERQSTSRRP..NTVOD..BASMRDVGGMNKPV-----	
ord	326	KRCFRDFCFPLKMRMERQSTSRRP..NTVOD..BASMRDVGGMNKPV-----	
AT1a	310	KKYELQLLKYIIPPKAKSHS...SLSTK..STLSYRPSDNMSSAKPASCFEVE-	
BK-2	340	RKKSWEVYQGVQKGGCRSEPIQOMENS..GTL..RTSISVEROIHKLQDWAGSRQ	

FIG. 14

28/31

SEQ ID NO: 83 mORmouse	1 MDSSAGPGNISDCSDPLA.PASCSPA..PGSWIWLSHVDGNCSDPCGPNRGLGGSHSLOC
SEQ ID NO: 79 mORrat	1 MDSSATGPGNISDCSDPLA.QASCSPA..PGSWIWLSHVDGNCSDPCGLNRGLGGNDSLC
SEQ ID NO: 84 mORbovin	1 MDSCAVPTNASNCIDPFTHPSCSPAPSPISSWVWNSHLÉGNLSDPCGPNRTELGCSDRLLC
SEQ ID NO: 85 mORhuman	1 MDSSAAPTNAASNCIDALAY.SSCSPAPSPGCVWVNLSHLDGNLSDPCGPNRDILGGNDSLC
SEQ ID NO: 86 mORpig	1 MDSSADPRNAASNCIDPESPSSMCSPVPSSWVNLSHLDGNLSDPCGPNRTELGCSDRLLC
SEQ ID NO: 87 mORws	1 METS...GNISDFLYPLS.....NSVMS.....NSSVLCRNFMSNTSFLNMNGSSRDSTD
SEQ ID NO: 81 ATla	1 -----MALNSSAEDGKRIQDDC
SEQ ID NO: 82 BK-2	1 -----MFSPWKISMFLSVREDSVPTTASFSADMNVTLQGPTLNG.TFAQSOK

mORmouse	58	POTGSPSPSMYTAITIMALYSIVCVVGLFGNLFVLMYVIVRYTKMKTATNIIYIFNLALADALA
mORrat	58	POTGSPSPSMYTAITIMALYSIVCVVGLFGNLFVLMYVIVRYTKMKTATNIIYIFNLALADALA
mORbovin	61	PSAGSPSPSMITAIIIMALYSIVCVVGLFGNLFVLMYVIVRYTKMKTATNIIYIFNLALADALA
mORhuman	60	FPTGSPSPSMITAIIIMALYSIVCVVGLFGNLFVLMYVIVRYTKMKTATNIIYIFNLALADALA
mORpig	61	FPTGSPSPSMITAIIIMALYSIVCVVGLFGNLFVLMYVIVRYTKMKTATNIIYIFNLALADALA
mORws	48	EODKICPVTIAITLTTLYSIVCVVGLFGNLFVLMYVIVRYTKMKTATNIIYIFNLALADALA
AT1a	19	RKACRHSYIIFVM..IPTLYSIIIFVVVGFTFGNSLVWIVIYFYMKIKITVASVFLNLALADLCF
BK-2	48	POVEWLGWLNTI..QPPFLWVFVLIATLEMI.FVLSVFCLHKSSCTVAEIIYLGNLAAADLIL

mORmouse	118	TSTLPPFQSVNLYLMG	TWPFGNLCKIVISIDYYNMFTSIFTLCTMSVDRYIAVCHPVKAL
mORrat	118	TSTLPPFQSVNLYLMG	TWPFGTILCKIVISIDYYNMFTSIFTLCTMSVDRYIAVCHPVKAL
mORbovin	121	TSTLPPFQSVNLYLMG	TWPFGTILCKIVISIDYYNMFTSIFTLCTMSVDRYIAVCHPVKAL
mORhuman	120	TSTLPPFQSVNLYLMG	TWPFGTILCKIVISIDYYNMFTSIFTLCTMSVDRYIAVCHPVKAL
mORpig	121	TSTLPPFQSVNLYLMG	TWPFGTILCKIVISIDYYNMFTSIFTLCTMSVDRYIAVCHPVKAL
mORws	107	TSTLPPFQSVNLYLMG	TWPFGDVKIVISIDYYNMFTSIFTLCTMSVDRYIAVCHPVKAL
AT1a	78	LLTLFLWAVTAMEYRWPFGNLCKIASSAASVTEINLYASVFFLTQPSIDRYLAIVHFMKSR	
BK-2	107	ACGLPPFWAITISNNFDWLFCTILCERVNIIISLMLYSSICFLMIVSLEDRYLAIVRTMSG	


mORmouse	177	DFRTPRNAAKIVNVCNWILSSAIGLPVMFMATTKYRQ	GSIDCTLTFSHPTWYWE
mORrat	177	DFRTPRNAAKIVNVCNWILSSAIGLPVMFMATTKYRQ	GSIDCTLTFSHPTWYWE
mORbovin	180	DFRTPRNAAKIVNVCNWILSSAIGLPVMFMATTKYRQ	GSIDCTLTFSHPTWYWE
mORhuman	179	DFRTPRNAAKIVNVCNWILSSAIGLPVMFMATTKYRQ	GSIDCTLTFSHPTWYWE
mORpig	180	DFRTPRNAAKIVNVCNWILSSAIGLPVMFMATTKYRQ	GSIDCALTFSHPTWYWE
mORws	166	DFRTPRNAAKIVNVCNWILSSAIGLPVMFMATTKYRQ	GSIDCTLTFSHPTWYWE
AT1a	138	LRRTPMLVAKIVTCIIWMLAGLSLPAVIRHVN	YFIENTNTIVCAFHESRNLTP
BK-2	167	RMRGVRWAKIVSLSVINGCILISSPMLVFRIMI	EYSDEGHNVTAQVISPM...LIWE

mORmouse	230	NLLKICVFI FAFIMPVL IIITCYGLMILRLKSVRMLSGSKEKDRNLRIRITRMVLVVVAVF
mORrat	230	NLLKICVFI FAFIMPVL IIITCYGLMILRLKSVRMLSGSKEKDRNLRIRITRMVLVVVAVF
mORbovin	233	NLLKICVFI FAFIMP [I IITCYGLMILRLKSVRMLSGSKEKDRNLRIRITRMVLVVVAVF
mORhuman	232	NLLKICVFI FAFIMPVL IIITCYGLMILRLKSVRMLSGSKEKDRNLRIRITRMVLVVVAVF
mORpig	233	NLLKICVFI FAFIMPVL IIITCYGLMILRLKSVRMLSGSKEKDRNLRIRITRMVLVVVAVF
mORws	226	TLLKICVFI FAFIMPVL IIITCYGLMILRLKSVRMLSGSKEKDRNLRIRITRMVLVVVAVF
AT1a	193	IIGLGTIKNLGFIFPFELIITSVTIWKALKIAYEIQKNNPRAADD...IEFRIMALVLF
BK-2	222	VETNMLLNVVGFLIP.LSIVTFCFTVQIMQWLRNNEQKFKIQTRE.RRATMVLVLLVLF

mORmouse	290	IVCWTPHIYVIKALITI	.	PETTFQTVSWHFCIALGYTNCLNPVLYAFLDENF
mORrat	290	IVCWTPHIYVIKALITI	.	PETTFQTVSWHFCIALGYTNCLNPVLYAFLDENF
mORbovin	293	IVCWTPHIYVIKALITI	.	PETTFQTVSWHFCIALGYTNCLNPVLYAFLDENF
mORhuman	292	IVCWTPHIYVIKALITI	.	PETTFQTVSWHFCIALGYTNCLNPVLYAFLDENF
mORpig	293	IVCWTPHIYVIKALITI	.	PETTFQTVSWHFCIALGYTNCLNPVLYAFLDENF
mORws	286	IVCWTPHIYVIKALITI	.	PNSTFQTVWHFCIALGYTNCLNPVLYAFLDENF
AT1a	250	FFSWTPHQIETFDVLQICGVIHDCKISDIVAPITLICLAVENCLNPI	FVFGFLGKKF	
BK-2	280	LICWLPFLQIESTFDLTHRGILSSCQDERIIDVITOIASEMYNSCLNPVLYVIVGKRF		

mORmouse	344	KRCFREFCG . I P T S S T I E Q Q N S A R I R Q N T R E H P S T A N T V D R T N H Q L E N L E A T A P L P
mORrat	344	KRCFREFCG . I P T S S T I E Q Q N S T R I R Q N T R E H P S T A N T V D R T N H Q L E N L E A T A P L P
mORbovin	347	KRCFREFCG . I P T S S T I E Q Q N S T R I R Q N T R E H P S T A N T V D R T N H Q L E N L E A T A P L P
mORhuman	346	KRCFREFCG . I P T S S N I E Q Q N S T R I R Q N T R E H P S T A N T V D R T N H Q L E N L E A T A P L P
mORpig	347	KRCFREFCG . I P T S S T I E Q Q N S A R I R Q N T R E H P S T A N T V D R T N H Q L E N L E A T A P L P
mORws	340	KRCFREFCG . V P S P S V L D Q N S T R N S M P Q C G O S S G H K V D R N A R O V
AT1a	310	K Y F L O L L K Y L P K A K S H S . . S L S T K M S T L S Y R S P D M S S A K P A C S C F E V E
BK-2	340	R K K S W E V Y O G V C O K G G C R S E P I O M E N S M G T L . R T S I S V E R O I H K L O D W A G S R O


FIG. 15



✓

30/31

FIG. 16 An Intracellular Point Mutation Results in Loss of Ligand-Induced Function



31/31

FIG. 17

